

## THE VIRGIN ISLANDS OF THE UNITED STATES

## VIRGIN ISLANDS DEPARTMENT OF HEALTH OFFICE OF THE COMMISSIONER

ST. CROIX OFFICE CHARLES HARWOOD COMPLEX 3500 ESTATE RICHMOND CHRISTINASTED, ST. CROIX, VI. 00820-4370 340-718-1311 ST. THOMAS OFFICE 1303 HOSPITAL GROUND, SUITE 10 CHARLOTTE AMALIE ST. THOMAS, VI 00802-6722 TEL: (340) 774-0117

September 9, 2019

To: All Parochial and Private Schools

Re: VACCINE REQUIREMENTS FOR SCHOOL ENTRY

(2019-2020 School Year)

The Virgin Islands Code Title 17, Chapter 10, Subchapter 116-1 states "all children registering for kindergarten and for any public, parochial or private schools must have a valid certificate of immunization for such diseases as Diphtheria, Pertussis, Tetanus, Measles, Poliomyelitis, and all other diseases as are required by the standard listing issued annually by the Commissioner of Health."

Title 17, Chapter 10, Subchapter 116-2 states "No principal or other person in charge of a kindergarten or other school shall knowingly admit or retain any pupil who has not submitted acceptable evidence of immunization according to the schedule specified by the Commissioner of Health, except even when there are exemptions as herein stated."

The following immunizations are <u>required</u> for school entry in the United States Virgin Islands for the school year 2019-2020:

Hepatitis B, Diphtheria, Tetanus, Pertussis, Polio, Measles, Mumps, Rubella, Varicella, and Meningococcal

- One (1) dose of Tetanus, diphtheria, acellular pertussis vaccine (Tdap) is required for children 11 years of age and above (eligibility for vaccination will be determined by health care provider).
- Two (2) doses of varicella vaccine are required for children in kindergarten through the 10<sup>th</sup> grade-both new and existing students who are entering/registering/repeating (eligibility for vaccination will be determined by health care provider).

- One (1) dose of Meningococcal conjugate vaccine (MCV4) for children 11 to 15 years of age, (eligibility to be determined by healthcare provider). A second dose of Meningococcal conjugate vaccine (MCV4) will be required for children 16 to 18 years of age, (eligibility to be determined by healthcare provider).
- Please note that in the 2019 -2020 school year, two (2) doses of varicella vaccine will be required for children in kindergarten through 11<sup>th</sup> grade.

For more information concerning the number of doses recommended based upon the child's age and immunization history, please refer to the attached 2019 "Recommended Immunization Schedule for Persons aged 0through 6 Years and Persons aged 7 to 18 years."

- Before being issued an immunization clearance form for school registration or being admitted to school, all children shall receive the required vaccine(s) (and total number of doses,) at the minimum age recommended in the attached immunization schedule; however, for each child, eligibility to receive the required vaccine will be determined by the clinic Nurse or healthcare provider.
- Any child who received the immunization clearance form at the time of the clinic visit but have
   <u>not</u> completed the required vaccines series ( due to the time intervals between doses, <u>must</u>
   return to the clinic or their healthcare provider to complete the required vaccine series by the
   vaccination date(s) indicated and documented by the immunization provider in the child's
   immunization record.

If you have any questions regarding the vaccines required for school entry, please call the Department of Health MCH Clinics in St. Thomas/St. John at (340) 777/8804/ St. Croix at (340) 514-6836.

Please note: Exemptions may be granted for school-aged children based on the principles and teachings of a recognized religious organization/church or if the vaccines/medicinal agents are medically contraindicated (please see VI Code Title 17, Chapter 10, Subchapter 116-3 &116-4).

Sincerely,

Justa Encarnacion

Date

Commissioner of Health

Pc: Principles, Administrators, and School Nurses

## Recommended Child and Adolescent Immunization Schedule for ages 18 years or younger United States, 2019 Table 1

These recommendations must be read with the Notes that follow. For those who fall behind or start late, provide catch-up vaccination at the earliest opportunity as indicated by the green bars in Table 1. To determine minimum intervals between doses, see the catch-up schedule (Table 2). School entry and adolescent vaccine age groups are shaded in grav.

| Vaccine                                                      | Birth    | 1 mo                      | 2 mos    | 4 mos                | 6 mos     | 80m 6                   | 12 mos                      | 15 mos                   | 18 mos                          | mos 2-3                                                     | 2-3 yrs   4-6                      | 4-6 yrs   7-1        | 7-10 yrs   11-1 | 11-12 yrs 13-15 yrs | yrs 16 yrs                     | 17-18 yrs         |
|--------------------------------------------------------------|----------|---------------------------|----------|----------------------|-----------|-------------------------|-----------------------------|--------------------------|---------------------------------|-------------------------------------------------------------|------------------------------------|----------------------|-----------------|---------------------|--------------------------------|-------------------|
| Hepatitis B (HepB)                                           | 1st dose | 2 <sup>nd</sup> dose      | ).<br>56 |                      |           |                         | . 3rd dose                  |                          |                                 |                                                             |                                    |                      |                 |                     |                                |                   |
| Rotavirus (RV) RV1 (2-dose<br>series); RV5 (3-dose series)   |          |                           | 1st dose | 2 <sup>nd</sup> dose | See Notes |                         |                             |                          |                                 |                                                             |                                    |                      |                 |                     |                                |                   |
| Diphtheria, tetanus, & acellular<br>pertussis (DTaP: <7 yrs) |          |                           | 1st dose | 2 <sup>nd</sup> dose | 3rd dose  |                         |                             | 4 <sup>th</sup> dose     | \$e                             |                                                             | <del>.</del> 5ф.                   | 5 <sup>th</sup> dose |                 |                     |                                |                   |
| Haemophilus influenzae type b<br>(Hib)                       |          |                           | 1st dose | 2 <sup>nd</sup> dose | See Notes |                         | 3rd or 4th dose. See Notes  | dose.⁴▶                  |                                 |                                                             |                                    |                      |                 |                     |                                |                   |
| Pneumococcal conjugate<br>(PCV13)                            |          |                           | 1st dose | 2 <sup>nd</sup> dose | 3rd dose  |                         | ▲4 <sup>th</sup> do         |                          |                                 |                                                             |                                    |                      |                 |                     |                                |                   |
| Inactivated poliovirus<br>(IPV: <18 yrs)                     |          |                           | 1st dose | 2 <sup>nd</sup> dose |           |                         | . 3 <sup>rd</sup> dose —    |                          | Î                               |                                                             | 4th (                              | 4 <sup>th</sup> dose |                 |                     |                                |                   |
| Influenza (IIV)                                              |          |                           |          |                      |           |                         | Anı                         | nual vaccina             | Annual vaccination 1 or 2 doses | loses                                                       |                                    |                      |                 | nnual vaccina       | Annual vaccination 1 dose only | nly               |
| Influenza (LAIV)                                             |          |                           |          |                      |           |                         |                             |                          |                                 | ٨                                                           | Annual vaccination<br>1 or 2 doses |                      | <b>3</b>        | nnual vaccina       | Annual vaccination 1 dose only | nly               |
| Measles, mumps, rubella (MMR)                                |          |                           |          |                      | See N     | Notes                   | <b>√</b> 1 <sup>st</sup> do | 1st dose▶                |                                 |                                                             | 2nd                                | 2nd dose             |                 |                     |                                |                   |
| Varicella (VAR)                                              |          |                           |          |                      |           |                         | 41st dose                   | <b>▼</b>                 |                                 |                                                             | 2nd (                              | 2 <sup>nd</sup> dose |                 |                     |                                |                   |
| Hepatitis A (HepA)                                           |          |                           |          |                      | See N     | Notes                   | 2-                          | 2-dose series, See Notes | See Notes                       |                                                             |                                    |                      |                 |                     |                                |                   |
| Meningococcal (MenACWY-D<br>≥9 mos; MenACWY-CRM ≥2 mos)      |          |                           |          |                      |           |                         | <b>4</b>                    | See Notes                |                                 |                                                             |                                    |                      | 1816            | 1st dose            | 2 <sup>nd</sup> dose           |                   |
| Tetanus, diphtheria, & acellular<br>pertussis (Tdap: ≥7 yrs) |          |                           |          |                      |           |                         |                             |                          |                                 |                                                             |                                    |                      | DT.             | Tdap                |                                |                   |
| Human papillomavirus (HPV)                                   |          |                           |          |                      |           |                         |                             |                          |                                 |                                                             |                                    |                      | S               | See<br>Notes        |                                |                   |
| Meningococcal B                                              |          |                           |          |                      |           |                         |                             |                          |                                 |                                                             |                                    |                      |                 | See                 | See Notes                      |                   |
| Pneumococcal polysaccharide<br>(PPSV23)                      |          |                           |          |                      |           |                         |                             |                          |                                 |                                                             |                                    |                      | See I           | See Notes           |                                |                   |
| Range of recommended ages for all children                   |          | Range of recommended ages | commende | dages                | œ .       | Range of recommended ag | Range of recommended ages   | ages                     | Range                           | Range of recommended ages for non-high-risk groups that may | ded ages fo                        | r non-high           | risk groups     | that may            | No reco                        | No recommendation |